THE IMPORTANCE OF DIFFERENT PARAMETERS OF GREATER SCIATIC NOTCH IN SEX DETERMINATION TILL THE AGE OF 20 YEARS

By

Adel A. EL-Hawary

From

Department of Anatomy, Faculty of medicine

Mansoura University, Mansoura, Egypt.

ABSTRACT

Various measurements of the greater sciatic notch (G.S.N.) e.g. width, depth and length of the posterior segment of the width were measured, and indices I and II were obtained from lateral radiographs of the pelvis of 168 subjects (78 males and 90 females). These subjects ranged in age from one to twenty years. Total and posterior angles were measured in each case.

Width, depth and posterior segment length of the G.S.N. were found valueless for sexing purposes before the age of 14 years. After that age till age of 20 years, these parameters become highly effective especially, in females, in sex determination. The length of the posterior segment of the width, posterior angle and index II were

found to be the best parameters, helping to assign the sex of the G.S.N. in different subjects before puberty. One can suggest that the growth of the width of the G.S.N. found in females before puberty occurs primarily in its posterior part.

INTRODUCTION

The importance of bones, in general, for sex determination is well documented both in anatomy (Boucher, 1957; Jit and Singh, 1966; Goldfarb, 1968) and anthropology (Davivongs, 1963). The greater sciatic notch (G.S.N.) attracted attention, as early as 1875, when Verneau (1875) observed that it was narrower in males and shallower in females. Jovanovic and Zivanovic (1965); Singh and Poturi (1978) and McMinn (1996) stated the influence of that sex on the G.S.N.

whereas the importance of G.S.N. in differentiation of the pelvic shape was denoted by Hollinshed and Rosse (1985); Llewellyn - Jones (1990); Lindner (1992) and Tortora (1995). Buchsbaum and Schmidt (1993) and Cunningham et al., (1994), also, emphasized the importance of the dimensions of the G.S.N in parturition.

Various measurements of the G.S.N. such as depth, length of the posterior segment and derived indices I and II in relation to sex determination were used (Williams et al., 1995). Davivongs (1963) found that female greater sciatic notches were deeper as well as wider in Australian aborigines. This deviation can be regarded as a racial characteristic. Width and depth of the G.S.N. may be significantly different in bones from different zones (Singh and Gangrade, 1968). These parameters were found to be valueless in sex determination (Singh and Potturi, 1978). Also, index I, which depends on the depth and width of the G.S.N. was accordingly not of much help in the sexing of hip bones. The posterior angle was the best single parameter, the length of the posterior segment and index 11 were high in females, as the widening

of the G.S.N. has occurred primarily in its posterior part (Singh and Potturi, 1978).

Most of the previous investigations were focused on the influence of the sex on the various parameters of the G.S.N. in adult human subjects. No attempt has been made to observe the influence of age (before puberty) on the different parameters of G.S.N. in both sexes. Therefore, the present investigation was undertaken to study the changes in parameters of the G.S.N. before puberty as they could be demonstrated radiologically and their relation to sex determination.

SUBJECTS AND METHODS

168 subjects were selected for this investigation (78 males and 90 females). They range in age from one to twenty years (Table 1). They were selected randomly from subjects referred to the Diagnostic Imaging Department, Mansoura Faculty of Medicine Hospital.

The subjects were clinically examined to exclude patients with skeletal deformities especially those of the spine and lower limbs. Other congenital or chronic bone diseases were also excluded. For each subject, a

plain x-ray of the pelvis in a true lateral position was done. In a perfect true lateral view film, the two greater sciatic notches were superimposed.

The following parameters of the G.S.N. were measured (Figs. 1-6) :

- 1 - The width of the G.S.N. (AB): the distance between the base of ischial spine (A) and the posterior inferior iliac spine (B).
- 2- The depth of the G.S.N. (CD): A line drawn from the deepest point of the concavity of the notch (C) perpendicular on the line AB, meeting it at the point (D)
- 3- The posterior segment of the width (BD).

$$4\text{- Index I : } \frac{\text{Depth (CD) x 100}}{\text{width (AB)}}$$

$$5\text{- Index II : } \frac{\text{Posterior segment x 100}}{\text{width (AB)}}$$

6- Total angle after construction on Radiography the triangle ABC and the depth CD, $\angle ACB$ denoted total angle.

7- Posterior angle: $\angle BCD$.

Statistical analysis :

Statistical analysis was performed using the student t-test paired values and p values of less than 0.05 were

considered significant (Armitage, 1983).

RESULTS

The principal parameters of the G.S.N. and their variabilities were studied in a sample of Egyptian population of both sexes at different ages from one to twenty years. The detailed observations are given in Tables 1,2 and in figures 1-13.

The depth of the G.S.N.:

In subjects before two years, the mean value of the depth of the G.S.N. was 2.55 ± 2.1 cm in male and 1.85 ± 3 cm in female. In both sexes this parameter showed gradual increase from the age of 6 years till the age of 20 years (Figure. 7). Before 10 years, the difference between the mean value of the depth in both sexes was not significant, although it was greater in males than in females. From 10-14 years, there was overlapping between the mean value of the depth in both sexes. The difference between the value of the depth in both sexes become significant ($p < 0.05$) from the age of 14 years onwards (Table 1 & Fig. 7).

The width of the greater sciatic notch (G.S.N.):

The width of the G.S.N. showed gradual increase in mean value in both sexes, as the age progresses till the age of 20 years (Fig. 8). In the young subject before 8 years, this measurement was found to be greater in males than in females. Between 8 to 14 years, it became greater in females than in males. These differences were not significant. From the age of fourteen years till the age of 18 years, the width of notch was found to be significantly greater in females than in males (Table 1 and Fig. 8).

Posterior segmen of the width of the G.S.N.:

Among the young subject before two years, the mean value of this parameter was greater in female ($1.1 \pm 0.3\text{cm}$) than in males ($0.8 \pm 0.4\text{cm}$). In males, as the age progressed, this parameter showed slowly and steadily increase in its value (Fig.9), where as in females, it showed rapid and progressive increase in values till the age of 20 years. The main value of the posterior segment was greater in females than in males at different ages. This difference was statistically significant ($p < 0.05$) between both sexes at different ages before 14 years, and it became highly significant ($p < 0.001$) after that

age (Table 1 and Fig. 9).

Index I of the G.S.N.:

Before two years, the mean value of index I was 55.8 ± 3.5 SD in males and 53.4 ± 7.1 SD in females. In the females, this index showed rapid and progressive decrease till the age of 20 years. In males, it showed progressive decrease till the age of 12 years. Thereafter, it increased rapidly and progressively till the age of 20 years. The index I of the notch was significantly

Index II of the G.S.N.:

In young subjects before two years the mean value of index II was 29.4 ± 3.8 SD in males and 42.8 ± 3.9 in females. In males this index decreased gradually in its value till the age of 18 years, whereas in females it decreased gradually in its value till the age of 12 years after which it showed rapid increase in its value till the age of 20 years. Index II in females was significantly greater than that of males at Jiff age erent ac groups (Table II & Fig. 11)

The total angle of the G.S.N.:

At different age groups, the total angle is greater in females than in males. Before 12 years, it seems to

be stationary in its value in both sexes, ranged between 81.7 to 82.5 in males and 87.9 to 86 in females. After that age, the value of this angle showed progressive and significant increase in females, and males till the age of 20 years. (Table 11 & Fig. 12)

The posterior angle of G. S. N.:
In young subjects before two

years, The man value of the posterior angle was 25.6 ± 7.2 in males and 36.5 ± 3.5 in females. In males, this angle decreased gradually in its value till the age of 20 years, whereas in females it showed gradual decreases in its value till the age of 20 years. Posterior angle value in females is significantly greater than that of males at different age groups. (Table II and Fig. 13).

Table (I) : The means and standard deviations of the various parameters of the greater sciatic notch (G.S.N.) (Depth, width, posterior segment and index I).

Age group (years)	No. of subjects		Depth		Width		Posterior segment		Index I	
	F.	M.	F.	M.	F.	M.	F.	M.	F.	M.
0-2 y	6	6	2.55 ± 2.1	1.85 ± 3.8	2.6 ± 4.2	2.7 ± 1.8	$1.1 \pm 3.6^*$	$0.8 \pm 3.4^*$	53.4 ± 7.1	55.8 ± 3.5
2-4 y	10	8	1.8 ± 3.2	1.6 ± 2.8	3.1 ± 2.6	3.4 ± 2.1	$1.25 \pm 4.5^*$	$0.9 \pm 4.1^*$	53.1 ± 7.3	54.3 ± 4.3
4-6 y	9	8	2.15 ± 4.2	1.45 ± 4.2	3.8 ± 1.8	3.4 ± 2.8	$1.3 \pm 4.1^*$	$0.95 \pm 2.6^*$	51.8 ± 2.6	54 ± 3.9
6-8 y	7	9	2.25 ± 0.2	2.05 ± 3.2	4.1 ± 1.6	4.2 ± 3.4	1.4 ± 3.1	$1.6 \pm 7.1^*$	51.4 ± 7.7	53 ± 3.6
8-10 y	8	6	2.45 ± 1.2	2.45 ± 3.1	4.8 ± 3.2	4.6 ± 2.8	$1.65 \pm 3.6^*$	$1.1 \pm 3.1^*$	50.8 ± 7.4	53.5 ± 4.1
10-12 y	10	8	2.6 ± 0.8	2.65 ± 3.1	5.4 ± 2.1	4.9 ± 1.8	$1.75 \pm 4.2^*$	$1.05 \pm 4^*$	50 ± 8.3	52.1 ± 4.9
21-14 y	7	8	2.8 ± 3.6	2.8 ± 3.6	5.6 ± 3.8	5.2 ± 1.8	1.95 ± 0.6	$1.05 \pm 3.1^*$	49.5 ± 7.5	54.2 ± 5.9
14-16 y	13	8	$3.1 \pm 1.2^*$	$2.9 \pm 1.8^*$	$6.0 \pm 5.1^*$	$5.3 \pm 4.1^*$	$2.3 \pm 3.1^{**}$	$1.15 \pm 4.5^{**}$	$48.3 \pm 10.3^*$	$58 \pm 7.9^*$
16-18 y	9	8	$3.2 \pm 4.2^*$	$2.85 \pm 1.1^*$	$6.1 \pm 3.1^*$	$5.4 \pm 3.2^*$	$2.4 \pm 2.8^{**}$	$1.1 \pm 2.1^{**}$	$47.9 \pm 2^*$	$59.6 \pm 7.3^*$
18-20 y	11	9	$3.25 \pm 2.1^*$	$2.8 \pm 3.5^*$	$6.2 \pm 4.2^*$	$6.6 \pm 3.8^*$	$2.5 \pm 1.8^{**}$	$1.15 \pm 2.8^{**}$	$46 \pm 4.1^*$	$59.9 \pm 4.9^*$
Total No	90	78								

F : Female M : Male

* Significant $P < 0.05$

** Highly significant $P < 0.001$

Table (II) : The means and standard deviations of the various parameters of the greater sciatic notch (G.S.N.) (index II, total angle and posterior angle) .

Age group (years)	No. of subjects		Index II		Total angle (in degrees)		Posterior angle (in degrees)	
	F.	M.	F.	M.	F.	M.	F.	M.
0-2 y	6	6	42.8±3.9	29.4±3.8*	85±4.5	82.5±4.5	36.5±3.5*	25.6±7.2*
2-4 y	10	8	38.6±3.8*	26.1±4.5*	84.8±4.4	83.8±4.6	36.5±4.1*	25.5±4.6*
4-6 y	9	8	34.9±4.6*	23.9±4.6*	85.5±5.5	83.2±5.8	36.5±4.4*	25.1±3.1*
6-8 y	7	9	±34.13.9*	24.1±4.5*	86±6.8	82.1±5.6	35.5±4.3*	23.3±4.1*
8-10 y	8	6	33.7±4.2*	23.9±5.1*	85±4.7	81.7±5.1	34.7±4.2*	22.6±5.2*
10-12 y	10	8	32.8±4.4*	21.2±3.8*	83.9±6.2	81.7±6.4	33.1±3.6*	21.7±4.8*
21-14 y	7	8	34.8±5.4*	21.1±3.4*	84±4.5	78.9±5.7	33.7±4.7*	21.5±5.1*
14-16 y	13	8	38.3±7.4**	20.1±4.5**	85.4±5.1*	76±6.2*	37.5±5.0**	19.7±3.6**
16-18 y	9	8	39.3±6.9**	19.5±3.9**	86.4±8.2*	72.1±8.2*	39.9±5.0**	18.2±3.4
18-20 y	11	9	39.9±5.9**	20.5±3.2**	88.5±8.1*	80.3±8.2*	41.2±4.1**	15.2±4.5**
Total No	90	78						

F : Female M : Male

* Significant $P < 0.05$

** Highly significant $P < 0.001$

Fig. 1 : Photography of true lateral view radiograph of the pelvis of male subject aged two years at the region of G.S.N

Fig. 2 : Photography of true lateral view radiograph of- the pelvis of female subject aged 3 years at the region of G.S.N

Fig. 3 : Photography of true lateral view radiograph of the pelvis of male subject aged 12 years at the region of G.S.N.

Fig 4 : Photography of true lateral view radiography of the pelvis of female subject aged 14 years at the region of G.S.N.

Fig. 5 : Photography of true lateral view radiography of the pelvis of male subject aged 18 years at the region of G.S.N.

Fig. 6 : Photography of true lateral view radiography of the pelvis of female subject aged 20 years at the region of G.S.N.

Fig. 7 : Depth of G.S.N. among males and females in different age groups.

Fig. 8 : Width of G.S.N. among males and females in different age groups.

Fig. 9 : Posterior segment of G.S.N. among males and females in different age groups.

Fig. 10 : Index I of G.S.N. among males and females in different age groups.

Fig. 11 : Index II of G.S.N. among males and females in different age groups.

Fig. 12 : Total angle of G.S.N. among males and females in different age groups.

Fig. 13 : Posterior angle of G.S.N. among males and females in different age groups.

DISCUSSION

For taking the different measurements of the G.S.N. two bony landmarks were selected: the posterior inferior iliac spine as the upper point and base of the ischial Spine as the lower point. This method was used by Litterman (1941). Davivongs (1963) and Singh and Potturi (1978) used the tubercle of the pyri formis as the upper point. Jovanovic and Zivanovic (1965) used the tip of ischial spine as lower point of notch instead of its base.

In the present study, both the width and depth of G.S.N. showed gradual increase in dimensions till the age of 14 years. The growth pattern of these parameters followed approximately the normal growth pattern of body (Myerscough, 1982). During this period, the greater sciatic notch was found to be wider in males. Male notches were found to be deeper. The differences of these parameters, in both sexes, were not significant. The present study showed that both width and depth are in fact of little value in sex determination during this period. Similar observations have been made earlier by Greulich and Thoms (1945) and Daivong (1963) who have reported

that both sexes start life with identical pelves and the major sexual differences appear at puberty. However, Boucher (1957) stated that the distinction between both sexes can be made even during foetal life.

Index I which depends on depth and width the G.S.N. was accordingly not of much help in the sexing of the G.S.N. before 14 years. However, it was found to be higher in males. This confirmed the earlier reports of Singh and Potturi (1978). From the age of fourteen years till 20 years, the greater sciatic notch was found to be significantly wider ($p < 0.05$) in females than in males, while male notches were found to be significantly deeper ($P < 0.05$) than those of females. Index I was accordingly found to be significantly higher in males. Similar observations were given before in adult subjects (Myerscough, 1982 and Lindner, 1992). Similarly, the total angle of the G.S.N. showed little differences between both sexes up to the age of fourteen years. After that, the total angle of the G.S.N. of the female pelvis become significantly greater than that of the male pelvis. Sexual discrimination of the G.S.N. could be made by the use of the posterior seg-

ment of the G.S.N., index II and posterior angle.

The fact that the parameters of the posterior segment and index II as good sex determination factor confirms the views of Jovanoic et al., (1968), who stressed the importance of the upper segment of the G.S.N in sex determination. The widening of the G.S.N found in females was found to be is an excellent parameter for discriminating sexual dimorphism at different age groups.

Singh and Potturi (1978) stated that, the adult pelvis, the width and depth of the G.S.N. are useless for determining the sex. The present study showed that beside the value of the depth, width and index I for sexing purposes in adult subjects, the parameters were useful for sex determination as early as fourteen years and to be useless criteria for sexing purpose before that age.

The length of the posterior segment of the width of the G.S.N, index II and posterior angle of the G.S.N was found to be the most useful parameters for sex determination at different ages before puberty. The mean

value of the posterior segment of the adult female were more two times those of male.

REFERENCES

- 1- Armitage, P. (1983) : Statistical methods in medical research. Blackwell scientific publications, Oxford, London vol. II, p. p. 168-172.
- 2- Boucher B. (1957) : Foetal sciatic notch; sex differences. Forensic Med., 1 : 51 -54.
- 3- Buchsbaum, H. and Schmidt, I.D. (1993) : Gynaecologic and obstetric surgery, 3rd. ed., Saunders W.B. Company, London, Philadelphia, Montreal, Tokyo, p.p. 25-28.
- 4- Cunningham, G. G.; MacDonald, P. C.; Leveno, k. J. et al., (1994) : Williams obstetrics. 19th. ed., Vol. 1, Elias Modern Press, London, p.p 290-294.
- 5- Davivongs, V. (1963) : The pelvic girdle of Australian aborigine; sex differences and sex determination Am. J.

Phys. Anthropol., 21: 444-455

- 6- Coldfarb A. F. (1968) : Dynamics and abnormalities of puberty. Clin. Obstet. Gynaecol., 11: 769-776.
- 7- Greulich, W. W. and Thoms, H. (1945) : The growth development of individual girls before, during and after puberty. J. Biol. Med., 17:91-97.
- 8- Hollinshead, W.H. and Rosse, C. (1985) : Textbook of Anatomy 4th ed, Harpet and Row publisher, Bhiladelphia; p.p. 737-740.
- 9- Jit, I. and Singh, S. (1966) : Sexing of adult clavicles; Indian J. Med. Res., 54: 551- 571.
- 10- Jovanovic S. and zivanovic, S. (1965) : The establishment of the sex by greater sciatic notch. Acta Anat. 61: 101-107.
- 11- Jovanovic, S.; Zivanovic, S. and Lotric, N. (1968) : The upper part of greater sciatic notch in sex determination of pathologically deformed hip bones. Acta Anat., 69: 229-238.
- 12- Litterman, G.S. (1941) : The greater sciatic notch in American Whites and Negroes. Am J. Physic. Anthropol., 28: 99-116.
- 13- Lindner, H. H. (1992) : Clinical Anatomy. Prentice - Hall international Inc., London, p.p. 458-460.
- 14- Llewellyn-Jones, D. (1990) : Fundamental of obstetrics and gynaecology vol . 1, ELBS wolfe Publisher, London, New York and Tokyo, p.p. 219-221.
- 15- McMinn, R. M. H. (1996) : Last's Anatomy, Regional and Applied, 9th ed., Churchill livingstone, Edinburgh London, New York and tokyo, p.p. 219-221 .
- 16- Myerscough, P.R. (1982) : Munro Kerr's operative obstetrics. 9th. ed., The English Language Book Society,

p.p. 38-46.

17- Singh, S. and gangrade, K. C. (1968) : Sexing of dult clavi-
cles: verification and appli-
cability of demarking points.
J. Ind. Acad. Forens. Sci., 7:
20-30.

18- Singh, S. and Potturi. B.R. (1978) : Greater sciatic noth
in sex determination. J.
Anat., 125: 619-624.

19- Tortora .J. (1995) : Principles of
Human Anat., Harper Col-
lins College Publisher, New
york, p.p. 162-164.

20- Verneau, R. (1875) : La bassin
dans les sexes et dans
les races. J. B. Bailliere
fils, paris, p. 63 (Cited
from: sing, S. and Pot-
turi, B.R. (1978) : Greater
sciatic notch in sex
determination. J. Anat, 125:
619-624.

21- Williams, P.L.; Bannister, L. H.; Berry, M. M., Collins, P. et al., (1995) : Morpho-
logical classification of
pelvis Gray's Anatomy
38th ed., Clurchill Living-
stone, London, p.p 670-
674.

أهمية القياسات المختلفة للشرم الوركى الكبير فى تحديد نوع الجنس حتى عمر العشرين عاماً

د. عادل عبد المهدى الهوارى

مدرس بقسم التشريح - كلية الطب - جامعة المنصورة

تم أخذ قياسات مختلفة للشرم الوركى الكبير مثل العرض والعمق والطول للجزء الخلفى من عرض الشرم الوركى وكذلك بعض المعاملات (مثل معامل ٢٠١) وقد أخذت هذه القياسات من مناظر الأشعة الجانبية للحوض فى ٧٨ ذكر و ٩٠ أنثى ولقد رتبنا هذه المقاسات حسب الأعمار المختلفة من سن عام حتى عشرون عاماً. ولقد تم قياس مجموع الزوايا وكذلك الزاوية الخلفية بعد عمل مثلث من المقاسات السابقة. ولقد وجد أن قياسات العرض والعمق والطول والمعامل الأول للشرم الوركى الكبير ليس لها قيمة لتحديد الجنس قبل سن الأربعة عشرة عام ولكن بعد هذا السن كانت هذه المقاييس ذات قيمة كبيرة وخاصة فى الإناث . ووجد أن طول الجزء الخلفى لعمق الشرم الوركى الكبير وكذلك زاوية الخلفية والمعامل الثانى هم أهم القياسات لتحديد نوع الجنس قبل سن البلوغ. ولقد أثبتت هذه الدراسة أن النمو فى عرض الشرم الوركى الكبير يتم فى الأنثى قبل سن البلوغ خاصة فى الجزء الخلفى.