INCIDENCE OF HDNB AND FOETOMATERNAL BLOOD GROUP INCOMPATIBILITY IN MANSOURA UNIVERSITY HOSPITALS

By

M. EL-ZINY and A. ABD EL-KADER

From

Pediatrics and Clinical Pathology Departments,

Faculty of Medicine, Mansoura University

Received for Publication : 3/4/1990

ABSTRACT

The study included 731 full term babies aged 1-6 days, presented with hyperbilirubinaemia (S.bilirubin > 12 mg/dl) in the period from 1985 to 1989. The babies and their mothers were subjected to 1) full history taking and thorough clinical examination 2) determination of S. bilirubin, Hb% and reticulocyte count in infants 3) blood group typing ABO, Rho and antibody screening for mothers and babies. In some instances they were typed for other blood group systems. 4) IgG antibodies (anti-A and anti-B) was done in ABO incompatible pregnancies.

It was found that out of the 731 cases with hyperbilirubinaemia, 73 babies (9.99%) had HDNB. In the later group, anti-D was detected in 24 infants (32.88%), anti-D and anti-C in 4 cases (5.48%), anti-c in 8 cases

(10.96%), ABO incompatibility in 34 cases (46.58%) and each of anti Fy^a, anti-Jk^a, & anti-K in one case (1.37%)

The results of the work showed that HDNB due to foetomaternal blood group incompatibility still a major problem which needs concern.

Abbreviation (Fy^a : Duffy a; Jk^a : Kidd a; K : Kell; D. C : direct coombs, E. T : exchange transfusion).

INTRODUCTION

Red blood cells contain blood group antigens inherited from the father that may be lacking in the mother. Due to foetomaternal haemorrhage, some mothers make blood group antibodies against these paternally derived antigen before and after birth. Maternal IgG antibodies are transported across the placenta and can cause haemolysis of foetal red blood cells.

MANSOURA MEDICAL JOURNAL

Haemolytic disease of newborn (HDNB) can vary in severity from hydrops foetalis in utero to only IgG sensitization of foetal red blood cells without haemolysis, (Synder et al., 1983, Quinn et al., 1988 and Ichikawa et al., 1989).

This work was intended to study

the incidence of HDNB in Mansoura University Hospitals due to foetomaternal blood group incompatibility in the period from 1985 to 1989. This

may throw a light on the importance of each blood group system incompatibility and help in selection of the suitable donor for exchange transfusion free from the accused antigen when needed. More over, we try to outline the approach to diagnosis in foetomaternal blood group incompatibility.

MATERIAL AND METHODS

This work was carried out on 731 full term babies aged 1-6 days presented to the neonatal intensive care unit with icterus and total serum bilirubin > 12 mg/dl. Each newborn was subjected to :

- 1- Full history taking and thorough clinical examination.

- 2- Laboratory investigations which

- 4- Direct antioglobulin test, heat elution of sensitized RBCs, then the eluted antibodies were tested using the commercial screening cells (Diamed cells).

Each mother of each selected newborn was subjected to :

- 1- Full history taking and details of delivery events.

- 2- Blood grouping, Rh typing and antibody screening (A. S) and in some instances they were typed for other systems. A. S was made by testing mother serum against two or three reagent red cell preparations "Diamed cell"

of known Phenotype using anti-globulin technique. The test procedures were selected to provide optimum conditions for detection of antibodies in the Rh, Kell, Duffy, Kidd, MNSs, Lewis, Lutheran & Xg systems.

When an antibody had been de-

tected, we identify it's specificity using (Diamed -Cell identification).

3-Indirect coombs test.

4- In ABO-incompatible pregnancies, antenatal tests for IgG antibodies (anti-A & anti-B) were introduced in vaccinated women and in those who experienced ABO-HDNB following prior delivery, IgM is neutralized with neutrab, or by agglutination with a panel of known group A and B and then separation of supernatant serum which were used in detection of IgG antibodies (Simmons, 1980).

RESULTS

The results of this work are shown in tables (1-6) and Fig. 1.

Table (I) shows the incidence of red cell antibodies causing the foetomaternal incompatibility among 731 cases of icteric newborn at Mansoura University Hospitals in the period from 1985 to 1989. From this table it could be observed that 73 cases (9.99%) are due to foetomaternal blood group incompatibility. The most predominant cause of HDNB is ABO

incompatibility (46.58%), then anti-D and anti-D plus anti-C accounting for about 38.36% of cases of HDNB, followed by anti-c (10.96%).

Tables (2-6) show the detailed results of foetomaternal incompatibility due to ABO blood grouping system, anti-D in d-women, anti-D in D and D^u-women, anti-D plus anti-C in ddcc and D^u cc-women and anti-c in CC - women respectively.

DISCUSSION

This work demonstrated that the most common cause of HDNB is ABO incompatibility with an incidence of 4.65% in relation to total icteric babies and 46.58% relative to type specificities. These data are moderate between the finding of Maisels and Giffords (1986) "20%" and that of Valentine (1958) "60%". Fourteen cases out of 34 were first born infants, this is in agreement with Mollison, (1979), Grundbacher (1980) and Haestis et al. (1988). Gupte and Bhatia (1980) explained the affection of first born infant by the formation of IgG antibodies by mothers on exposure to various agents e.g. food stuffs, infective agents and vaccines which contain A or B like substances. As regard the blood grouping, we found O-

A combination in 20 situations and O-B combination in 14 (Table 2). This goes with the finding of Bengtsson and Virnehoit 1974. DCT was found to be negative in 35.25% of cases with ABO incompatibility, which goes with the finding of Oski (1984) and Huestis et al., (1988).

However, antibody detection in elute in DCT negative babies was found to be positive in 10 cases out of 12 and IDC test for mothers was found to be positive. So in suspected ABO-HDNB with negative DCT we must do IDC test for mothers and antibody detection for the elute of red cells of the infant. Moreover, IgG (anti-A and anti-B) was found to be more than 64 mg/dl in all mothers with ABO incompatibility.

The Lw gene is regulated by the Rh gene, its absence lead to absence of Lw antigen inspite of presence of Rh-gene. The majority of Rho positive cells are Lw-positive, so anti-LW reacts with almost all Rho Positive cells, in this case anti-LW as anti-D is suggested (Tippett, 1977).

HDNB caused by anti-c was found in 10.96% of cases (Tables 1 and 5). an incidence agree with that recorded by Smith et al. (1967) and Hardy and Napier (1981). This finding denote the importance of phenotyping for complete Rhesus blood grouping of young girls and women before blood transfusion to avoid immunization by supplying blood lacking c-antigen for those negative for this antigen. Phenotyping of red cells of babies is also essential for confirming the allo immunization.

Moreover, HDNB could be caused by other blood group systems than Rh and ABO as anti-K, anti-Fy^a and anti-JK^a (Soleta et al., 1983).

Caine and Heubach (1986) reported that anti-K is the most potent among the irregular antibodies; after those of Rh system, which can cause severe HDNB and still-birth. So we advocate the use of only K-negative blood for female recipients.

Antibodies of the Duffy system come next to those of Rh and Kell systems among the irregular antibodies causing HDNB (Shah & Gilyas, 1983). Duffy sensitization must be managed in a manner similar to Rh sensitization, with amniocentesis being performed at the appropriate time as it is diagnostic and is of higher predictability (Sturket et al., 1988). The blood bank should be notified prior to delivery of a possible affected foetus, so that Fy(a-b-) or Fy(a-B+) blood can be obtained for exchange transfusion if indicated (Contreras et al., 1983).

HDNB caused by anti-JK^a is mentioned by many investigators (Simpson et al., 1978, and Harrison

and Popper, 1981).

In this work, each of anti-K, anti-Fy^a and anti JK^a was found in one case (1.37%), and this goes with the Previous reports.

CONCLUSION

- ABO incompatibility is the most common cause of HDNB in our locality (46.58%) and the best Predictor for ABO-HDNB is the titre of maternal IgG anti-A or anti-B, and the results of direct antiglobulin test on RBCs of cord blood fortified by elute examination for antibodies in cases with negative D.C.

- Rh D-mediated HDNB still significant (38.36%), though Rh-immunoglobulin prophylaxis was introduced since many years, so it needs reevaluation. Also, the use of anti-D immunoprophylaxis in DU variants is suggested.

- HDNB can occur secondary to foetomaternal incompatibility in minor blood group systems so it is important to select the donor lacking the corresponding antigen for ET of the affected infant.

Table (1) : Incidence of red cell anti - bodies causing haemolytic disease among 731 cases of icteric newborn in Mansoura University Hospital.

Cause	NO. of Cause	Frequency in relation to total specimens (%)	Frequency in relation to HDNB (%)
- Anti - D	24	3.28	32.88
- Anti - D +	4	0.55	5.48
Anti - C			
- Anti - c	8	1.09	10.96
- ABO	24	4.65	46.58
incompa - tibility			
- Anti - Fy ^a	1	0.14	1.37
- Anti - JK ^a	1	0.14	1.37
- Anti - K	1	0.14	1.37
Total	73	9.99	

Table (2) : HDNB due to ABO incompatibility .

	NO . of cases	% relat . to all Icteric Infants	% re lat. to HDNB	% relat . to ABO Incompa - tibility	Remarks
Total NO .	34	4.65	46.58	100%	- 2 infants recover without treatment
Blood group Combinations					
O - A	20	2.74	27.4	58.82	- 11 treated with pho -
O - B	14	1.92	19.9	41.18	totherapy
					- 21 treated with pho -
DCT of cord blood					therapy & exchange transfusion
- Positive	22	3.02	30.14	64.75	
- Negative	12	1.63	16.44	35.25	
E lute in negative DC	10	1.37	13.7	29.41	
NO . of mothers with positive IDC (Ig G antibody > 64)	32	4.38	43.12	43.12	

Table (3) : pregnancy outcome in d - women with D - husband and significant Anti - D .

Cause	group & pheno	Maternal	Antecedent risk Fs	Cord blood	D.C.	Remarks
NO .	type	partly	HDNB	Abortion	Bl, trasl	-----
1	A 1 B - ccdee	2nd	-	-	-	A 1 B - ccDee
2	A 1 ccdee	2nd	-	-	-	O - ccDEE
3	A 1 ccdee	2nd	1st	one	-	B - CcDee
4	O - ccdee	2nd	-	-	-	O - CcDee
5	AB - ccdee	4th	2nd &	-	-	A 1 - CcDee
6	O - ccdee	4th	2nd	-	-	O - ccDee
7	A 1 B - ccdee	3rd	2nd	-	-	A 1 - ccDee
8	B - ccdee	3rd	-	-	-	O - ccDee
9	O - ccdee	4th	2nd &	-	-	O - ccDee
10	O - ccdee	3rd	2nd	-	+	O - ccDee
11	A 1 B - ccdee	3rd	2nd	-	-	A 1 ccDee
12	B - ccdee	3rd	-	-	-	O - ccDee
13	O - ccdee	4th	2nd	-	-	A 1 - CcDee
14	A 1 B - ccdee	3rd	-	-	-	B - ccDee
15	B 1 ccdee	4th	-	-	-	B, ccDee
16	B, ccdee	2nd	-	one	-	O, ccDee
17	A 1 B, ccdee	4th	2nd &	3rd	-	A 1, ccDee

ET : Exchange trans fusion

Table (4) : pregnancy outcome in D and Duwouen with significant anti - D and D - husband .

case No .	Maternal blood group and pheno type	Antecedent risk factors	Cord blood group	D . C .	Re marks
1	A 1 B - ccD ^U Ee	6 th para - from 2 nd to 5 th infant had HDNB & died .	B - ccDEe	+++	Died
2	A 1 - ccD ^U ee	3 rd para	perinatal death	-	
3	O - ccD ^U ee	2 nd para	O - Cc Dee	++	ET
4	A 2 B - ccD ^U ee	3 rd gravida , 2 nd 1 abour with HDNB	A2 - ccDee	+	ET
5	A 1 - ccD ^U ee	2 nd pare	A 1 - ccDee	+	ET
6	A 1 - ccDee	6 th para - previous 5 infants with good conditions .	O - CcDee	+++	ET
7	B - ccDee	5 th para - previous still birth	B - ccDee	++	ET

Table (5) : pregnancy outcome with anti - D + anti C

case No.	Maternal blood group and pheno type	Antecedent risk factors	Cord blood group	D . C . Re marks
1	A 1 B - ccdee	8 th para, 2 nd to 7 th infants died with HDNB	perinatal death	--
2	B - ccD ^u Ee	6 th para 5 th died with HDNB	O - Cc DEe	+++ one ET & died
3	A 1 B - ccD ^u ee	5 th gravida with history of previous HDNB	B - CcDee	++ one ET & died
4	B - ccD ^u Ee	7 th gravida with history of multiple HDNB	B - CcDee	++ two ET & died

Table (6) : pregnancy outcome in patients with anti - c

case No .	Maternal blood group and pheno type	Antecedent risk factors	Cord blood group	D . C .	Re marks
1	A 1 CCDee	2 nd para	B - ccDee	++	1 ET
2	B - CCDee	3 rd para	O - CcDee	+++	2 ET
3	B - CCDee	4 th para with parinatal death of 2nd infant	B - CcDee	+++	2 ET
4	O - CCDEe	3 rd para	perinatal death		
5	O - CCDee	4 th para , 2 nd & 3 rd child died with HDNB	O - CcDee	++	
6	O - CCDee	6 th para	B - Cc DEe	++	1 ET
7	O - CCDEe	2 nd para	B - CcDEe	+++	1 ET
8	A 1 - CCDEe	2 nd para	A 1 - CcDEe	++	1 ET

Fig. 1
Incidence of red cell anti-bodies causing HDNB among 731 jaundiced newborns

REFERENCES

- Bengtason. B and Verneholt, J.
(1974) : *Acta. Ped.*
Scand. 63 : 70.
- Caine. M. E and Heubach, E. M.
(1986) : *Am. J. Obstet. Gynaec.* 154 : 85.
- Clarke, C. A. : Mollison, P. L. and Whitfield, A.G.W.
(1985) : *Brit. Med. J.*
- Contreras, M.; Gordon, H. and Tidmarsh, E. (1983) :
Brit. J. Haemat. 53 : 355.
- Grundbacher, F. J. (1980) : *Transfusion* 20 : 563.
- Gupte, S. C. and Bhatia, H. M.
(1980) : *Vox. Sang.* 38 : 22.
- Hardy J. and Napier, J. A. F.
(1981) : *Brit. J. Obstet. Gynaec.* 88 : 9.
- Harrison, J. K. and Popper, E. L.
(1981) : *Transfusion* 21 : 90.
- Huestis. D. W.; Bove, J. R. and Case, I. (1988) : I) Practical blood transfusion (4th ed) Chap. 2, little Brown & Co. Bosion, Toronto 398.
- Ichikawa, Y.; Sato, C.; Mc Creary, J. and Lubenko, A.
(1989) : *Vox. Sang.* 56 : 98.
- Lacey, P. A.; Caskey, C. R.; Werner, D. J.; Moulds. J. J. 1983) : *Transfusion* 23 : 91.
- Maisels, M. J. and Gifford, K.
(1986) : *Pediatrics* 78 : 837.
- Mollison, P. L. (1979) : *Blood transfusion in clinical medicine* (6th ed) Chap. 16 Oxford Blackwell Scientific Publication 666.
- Oski, F. A. (1984) : Jaundice, 1. In Avery M. E.; Taeusch, H. W. (eds). Chapter 10. *Shaffer's Disease of the Newborn* W. B. Saunders Co. 622.
- Ostgard, P.; Eevang, F. and Kornstad, L. (1986) : *Acta. Pediat. Scand.*

- nisse Zentralbl. Gyna-
kol. 110 (1) :12.
- Synder, E. L.: Kennedy, M. S.:
Blumberg, N.: Bo-
rucki, D. T.: Lichtiger,
B.; Mc Carthy, L.;
Menitove, J. and Ol-
ive, J. A. (1983) :
Blood transfusion
therapy. American as-
sociation of blood
banks. Joyce Duncan,
Publications, Mango
Printed in the USA.
PP. 45.
- Tippett, P. and Sanger, R. (1977) :
Arzt. L. Lab. 23 : 476.
- Values, T. and Hyte, M. (1977) :
Pediatrics 59: 881.
- Valentine, G. H. (1958) : Arch. Dis.
Child. 33 : 185,
- White, C. A.; Stedman, C. M. and
Frank, S. (1983) : Am.
J. Obstet. Gynec. 145
: 1069.
- Wilken, H. (1988) : Zentralbl. Gy-
nakol. 110 : 1.
- Quinn, M. W.: Weindling, A. M. and
Davidson, D. C (1988)
: Arch. Dis. Child. 63 :
1258.
- Shah V. P. and Gillja, B. K. (1983) :
Nystate. J med. 2 :
244.
- Simmons, A. (1980) : Technical
haematology 3rd ed.
J. B. Lippincott Com-
pany. Philadelphia-
Tornoto P. 471.
- Simpson, M. B.: Pryzblylik, J. A.;
Denham, M. A. and
Rad Cliffe, J. H.
(1978) : Obst. Gynec.
52 : 616.
- Smith, B. D.: Harber, J. M. and
Gyeenan, J. T. (1967)
: Obstet. Gynaec. 29 :
119.
- Soleta, A.; Sibai, B. and Masen, J.
(1983) : Obstet. GY-
naec. 61: 25.
- Sturk, K.: Scharzlos, G. and Dur-
ing, R. (1988): Ergeb-

الملخص العربي

معدل حدوث مرض تكسير الدم عند الأطفال حديثي الولادة وعدم توافق فصائل الدم بين الطفل والأم في مستشفيات جامعة المنصورة

أجرى هذا البحث في الفترة من ١٩٨٥-١٩٨٩ وشمل ٧٣١ طفلاً كامل النمو يتراوح عمره بين ١-٦ أيام ويشكو من اليرقان، وكانت نسبة البيلروبين أكثر من ١٢ مجم لكل ١٠٠ سم^٣ من مصل الدم.

وقد تم عمل الآتي لكل من الأم والطفل :

(١) التاريخ المرضي والفحص الكلينيكي الدقيق.

(٢) تحديد نسبة البيلروبين في مصل الدم ونسبة الهيموجلوبين وعد الخلايا الشبكية في دم الطفل.

(٣) تحديد فصائل الدم (ABO) ، (Rh) وفحص الدم بحثاً عن الأجسام المضادة عند الأطفال والأمهات وتحديد فصائل الدم الأخرى في الحالات الإيجابية.

(٤) تحديد كمية الأجسام المضادة (ج) المضاد لنوع الفصيلة ،

وقد خلص البحث لما يأتي :

(١) عدد ٧٣ طفلاً كانوا مصابين بمرض تكسير الدم عند الأطفال حديثي الولادة نتيجة عدم توافق دم الطفل مع الأم بنسبة ٩٩.٩٪ من حالات اليرقان المبكر.

(٢) الأجسام المضادة لـ (Rh D) كانت موجودة في ٢٤ طفلاً من الأطفال حديثي الولادة المصابين بتكسير في الدم بنسبة ٣٢.٨٨٪ من هؤلاء الأطفال، والأجسام المضادة لـ Rh C + Rh D في ٤ حالات ٥.٤٨٪ والأجسام المضادة لـ Rh C في ٨ حالات (١.٠٩٨٪).

(٣) مرض تكسير الدم بسبب عدم التوافق في فصيلة الدم (ABO) كان موجوداً في ٣٤ حالة بنسبة ٤.٦٥٨٪.

(٤) الأجسام المضادة لكل من K, Jk^a, Fy^a كل منهما كان موجوداً في حالة واحدة بنسبة ١.٣٧٪ من حالات تكسير الدم في الأطفال حديثي الولادة.

ومن هذا يتضح أن مرض تكسير الدم عند الأطفال حديثي الولادة بسبب عدم توافق دم الطفل مع الأم مازال يمثل مشكلة تحتاج لمزيد من الاهتمام.

TO THE SECRETARY OF THE ARMY
FROM THE SECRETARY OF THE ARMY

THE SECRETARY OF THE ARMY
THE SECRETARY OF THE ARMY

THE SECRETARY OF THE ARMY
THE SECRETARY OF THE ARMY

THE SECRETARY OF THE ARMY
THE SECRETARY OF THE ARMY

THE SECRETARY OF THE ARMY
THE SECRETARY OF THE ARMY

THE SECRETARY OF THE ARMY
THE SECRETARY OF THE ARMY

THE SECRETARY OF THE ARMY
THE SECRETARY OF THE ARMY

THE SECRETARY OF THE ARMY
THE SECRETARY OF THE ARMY

THE SECRETARY OF THE ARMY
THE SECRETARY OF THE ARMY

THE SECRETARY OF THE ARMY
THE SECRETARY OF THE ARMY

THE SECRETARY OF THE ARMY
THE SECRETARY OF THE ARMY

THE SECRETARY OF THE ARMY
THE SECRETARY OF THE ARMY

THE SECRETARY OF THE ARMY
THE SECRETARY OF THE ARMY

THE SECRETARY OF THE ARMY
THE SECRETARY OF THE ARMY

THE SECRETARY OF THE ARMY
THE SECRETARY OF THE ARMY

THE SECRETARY OF THE ARMY
THE SECRETARY OF THE ARMY